

On factorization of operators between Banach spaces

Eugene Tokarev

ABSTRACT. A finite-dimensional analogue $GL_{fin}(X)$ of the known Gordon-Lewis constant $GL(X)$ of a Banach space X is introduced; in its definition are used only finite rank operators. It is shown that there exist Banach spaces X such that $GL_{fin}(X)$ is finite and $GL(X)$ is infinite. Moreover, for any Banach space X of cotype 2, the finite-dimensional constant $GL_{fin}(X) < \infty$.

1. Introduction

Let X, Y be Banach spaces; $u : X \rightarrow Y$ be a (linear, bounded) operator. This operator is said to be L_1 -factorable if there exists a measure μ and operators $v : X \rightarrow L_1(\mu)$; $w : L_1(\mu) \rightarrow Y^{**}$ such that $k_Y \circ u = v \circ w$, where k_Y is a canonical embedding of Y into Y^{**} . It is clear that for any L_1 -factorable operator u

$$\gamma_1(u) := \inf\{\|v\| \|w\| : k_Y \circ u = v \circ w\} < \infty,$$

where the infimum is taken over all possible factorizations of u .

Grothendieck asked ([1], problem 2, p. 78):

Whether there exists absolutely summing operators which are not L_1 -factorable?

Gordon and Lewis [2] answered this question in affirmative. At the same time they showed that there exists a wide class of Banach spaces X with the following property:

(GL) *Every absolutely summing operator $u : X \rightarrow Y$ where Y is an arbitrary Banach space is L_1 -factorable.*

This class contains all *spaces with the local unconditional structure* (shortly, l.u.st.), which were introduced in [2]. Recall that according to [3] a Banach space X is said to have the l.u.st. if and only if its second conjugate is isomorphic to a complemented subspace of some Banach lattice. Particularly, the class of all Banach spaces having the l.u.st. contains all Banach lattices. The result of [2] further was improved (see e.g. [4], p. 107); now it is known that if a Banach space X is of cotype 2 and has the l.u.st. then each its subspace enjoys the property (GL).

Gordon and Lewis [2] defined for a given Banach space X a constant

$$GL(X) = \sup\{\gamma_1(u) / \pi_1(u) : u : X \rightarrow Y\},$$

where the supremum is taken over all Banach spaces Y and all absolutely summing operators $u : X \rightarrow Y$ (here and below $\pi_1(u)$ denotes the absolutely summing norm of an operator u). In their solution of the aforementioned Grothendieck problem

1991 *Mathematics Subject Classification.* Primary 47A68, 47B10; Secondary 46B07, 46B20.
Key words and phrases. Banach spaces, Linear operators, Factorization.

a sequence $(u_n)_{n < \infty}$ of finite rank operators $(u_n : X \rightarrow l_2^{(m_n)})$ was constructed in such a way that $\sup\{\gamma_1(u_n)/\pi_1(u_n) : n < \infty\} = \infty$.

Let us define a *finite dimensional version of the Gordon-Lewis constant*:

$$GL_{fin}(X) = \sup\{\gamma_1(u)/\pi_1(u) : u : X \rightarrow l_2; \dim u(X) < \infty\}.$$

It is clear that $GL_{fin}(X) = \infty$ implies that $GL(X) = \infty$.

However in a general case may be a situation when for a given Banach space X constants $GL_{fin}(X)$ and $GL(X)$ are essentially different. To be more exact, in this article will be shown that for some Banach spaces X the constant $GL_{fin}(X)$ is finite and $GL(X)$ is infinite.

The question of the full description of a class **GL** of those Banach spaces that has the property (GL) (i.e. those Banach spaces X for which $GL(X) < \infty$) is open. In the article it would be shown that a class of all Banach spaces X for which $GL_{fin}(X) < \infty$ contains the class of all Banach spaces of cotype 2.

2. Definitions and notations

Let $\mathcal{B}$ denotes a class of *all* Banach spaces. For $X, Y \in \mathcal{B}$ let $B(X, Y)$ be a Banach space of all bounded linear operators $u : X \rightarrow Y$ (equipped with a strong operator topology).

Let $1 \leq p \leq \infty$. An operator $u \in B(X, Y)$ is said to be

- *p-absolutely summing*, if there is a constant $\lambda > 0$ such that

$$(\sum_{j < n} \|u(x_j)\|^p)^{1/p} \leq \lambda (\sum_{j < n} |\langle x_j, f \rangle|^p)^{1/p}$$

for any $f \in X^*$ and any finite set $\{x_i : i < n; n < \infty\} \subset X$.

Its *p-absolutely summing norm* $\pi_p(u)$ is the smallest constant λ .

- *p-factorable*, if there exists a such measure μ and operators $v \in B(X, L_p(\mu)); w \in B(L_p(\mu), Y^{**})$ such that $k_Y \circ u = w \circ v$, where k_Y is the canonical embedding of Y in its second conjugate Y^{**} , and the symbol $\circ$ denotes the composition of operators.

Its *γ_p -norm* is given by

$$\gamma_p(u) = \inf\{\|v\| \|w\| : k_Y \circ u = w \circ v\}.$$

- *p-integral*, if there exists a such probability measure μ and such operators $v \in B(X, L_\infty(\mu)), w \in B(L_p(\mu), Y^{**})$ that $k_Y \circ u = w \circ \varphi \circ v$, where φ is an inclusion of $L_\infty(\mu)$ into $L_p(\mu)$.

Its *p-integral norm* is given by

$$\iota_p(u) = \inf\{\|v\| \|\varphi\| \|w\| : k_Y \circ u = w \circ \varphi \circ v\}.$$

- *Nuclear*, if there are sequences $(f_n)_{n < \infty} \in X^*; (y_n)_{n < \infty} \in Y$ such that for all $x \in X$

$$u(x) = \sum_{n < \infty} f_n(x) y_n.$$

Its *nuclear norm* $\nu_1(u)$ is given by

$$\nu_1(u) = \inf\{\sum \|f_n\| \|y_n\| : u(x) = \sum_{n < \infty} f_n(x) y_n\},$$

where the infimum is taken over all possible representations of u .

A Banach space X is said to be of *cotype 2* if there exists such a constant $c < \infty$ that for any $n < \infty$ and any subset $\{x_i : i < n\} \subset X$

$$\left(\sum_{i < n} \|x_i\|^2 \right)^{1/2} \leq c \int_0^1 \left\| \sum_{i < n} r_k(t) x_k \right\| dt,$$

where $r_k(t)$ are Rademacher functions.

It is well known that the space L_p is of cotype 2 for all $p \in [1, 2]$.

3. Main results

DEFINITION 1. Let Y be a Banach space of cotype 2. It will be said that $X \in \mathcal{B}$ is the Pisier's space for Y , if following conditions are satisfy.

1. Y is isometric to a subspace of X . X and X^* both are of cotype 2.
2. Any operator $u : X \rightarrow l_2$ is absolutely summing; any operator $v : X^* \rightarrow l_2$ is absolutely summing.
3. There is a constant λ_X such that for any finite rank operator $u : X \rightarrow X$ its nuclear norm $\nu_1(u)$ satisfies the estimate $\nu_1(u) \leq \lambda_X \|u\|$.
4. If X contains a subspace isomorphic to l_1 then its conjugate X^* contains a subspace isomorphic to l_2 .
5. Any operator $w : l_2 \rightarrow X$ is 2-absolutely summing; the same is true for every operator $s : l_2 \rightarrow X^*$.

The famous result of J. Pisier [5] shows the existence of the corresponding Pisier's space X for any Banach space Y of cotype 2.

THEOREM 1. There exists a Banach space X such that $GL(X) = \infty$ and $GL_{fin}(X) < \infty$.

PROOF. Let $X \in \mathcal{B}$ be a space which is the Pisier's space for $L_1[0, 1]$. Certainly, it has subspaces that are isometric to l_1 and to l_2 . Let l_2 be isometric to a subspace $X_2 \hookrightarrow X$ and $i : l_2 \rightarrow X_2$ be the corresponding isometry. Recall that X is of cotype 2 and that each operator $u \in B(X, l_2)$ is absolutely summing. Let $u : X \rightarrow l_2$. The composition $w = i \circ u$ acts on X , is absolutely summing and is L_1 -factorable if and only if u is. Assume that u has a finite dimensional range, i.e. that $\dim(u(X)) < \infty$. Then w is also a finite rank operator. Hence, its nuclear norm $\nu(w)$ may be estimated as $\nu(w) \leq \lambda \|w\|$, where λ does not depend on w . Thus,

$$\gamma_1(u) = \gamma_1(w) \leq \iota_1(w) \leq \nu(w) \leq \lambda \|w\| \leq \lambda \pi_1(w).$$

This sequence of inequalities implies that

$$\gamma_1(u) \leq \lambda \pi_1(u).$$

Recall that u was an arbitrary operator of finite dimensional range. Hence, $GL_{fin}(X) < \infty$.

Let us show that for a special choosing of $u : X \rightarrow l_2$ such inequalities may be not longer true. Indeed, according to [5], if X contains a subspace isomorphic to l_1 then its conjugate X^* contains a subspace (say, Y_2) which is isomorphic to l_2 . Let $j : l_2 \rightarrow Y_2$ be the corresponding isomorphism; $\|j\| \|j^{-1}\| < \infty$. Because of reflexivity, Y_2 is weakly* closed and, hence, there exists a quotient map $h : X \rightarrow (Y_2)^*$ which generates a surjection $H : X \rightarrow l_2$: $H = (j_2)^* \circ h$. From our assumptions on properties of X follows that the conjugate operator H^* is L_∞ -factorable and is 2-absolutely summing. Clearly, H^* induces an isomorphism between l_2 and $H^*(l_2)$.

However this property contradicts with $\pi_2(H^*) < \infty$. Hence, H cannot be factored through $L_1(\mu)$. $\square$

THEOREM 2. *For every Banach space of cotype 2, $GL_{fin}(X)$ is finite.*

PROOF. Let a Banach space Y is of cotype 2. It can be isometrically embedded in a corresponding Pisier's space X_Y , which has the same properties as was listed above. Let $T : Y \rightarrow l_2$ be an absolutely summing operator. Then T is 2-absolutely summing and, consequently, 2-integral, i.e. admits a factorizations $T = k \circ j \circ s$, where $s : X \rightarrow L_\infty(\mu)$; $k \circ j : L_\infty(\mu) \rightarrow l_2$. Because of injectivity of $L_\infty(\mu)$, s can be extended to $\bar{s} : X_Y \rightarrow L_\infty(\mu)$. The corresponding operator $\bar{T} = k \circ j \circ \bar{s} : X_Y \rightarrow l_2$ is 2-absolutely summing too and extends T . Recall that from properties of X_Y follows that $\bar{T}$ is absolutely summing. Assume that T has a finite dimensional range. The same is true for $\bar{T}$. From theorem 1 follows that $\bar{T}$ is L_1 -factorable. Hence T is also L_1 -factorable. $\square$

REMARK 1. *From [6] follows that if there exists some $p > 2$ such that the space l_p is finitely representable in $X \in \mathcal{B}$ then X contains a subspace Z for which $GL_{fin}(Z)$ is infinite.*

PROBLEM 1. *Whether every Banach space Z which is not of cotype 2 contains a subspace which does not enjoy this property?*

ACKNOWLEDGEMENT 1. *Author wish to indebted his thanks to V.M. Kadets for his attention to this paper*

4. References

1. A. Grothendieck, *Résumé de la théorie métrique des produits tensoriels topologiques*, Bol. Soc. Math. São Paulo, **8** (1956), 1-79
2. Y. Gordon and D.R. Lewis, *Absolutely summing operators and local unconditional structures*, Acta Math., **133** (1974), 27-48
3. T. Figiel, W.B. Johnson and L. Tzafriri, *On Banach lattices and spaces having local unconditional structure with applications to Lorentz function spaces*, J. Approx. Theory, **13:4** (1975), 392-412
4. J. Pisier, *Factorization of linear operators and geometry of Banach spaces*, Publ.: Conf. Board of Math. Sci. by Amer. Math. Soc., Providence, Rhode Island, 1986, 153 p.
5. J. Pisier, *Counterexample to a conjecture of Grothendieck*, -Acta Math. **151** (1983) 181-208
6. T. Figiel, S. Kwapien and A. Pełczyński, *Sharp estimates for the constants of unconditional structure of Minkowski spaces*, Bull. Acad. Pol. Sci., Sér. Math. Astron. et Phys., **25** (1977), 1221-1226

B.E. UKRECOLAN, 33-81, ISKRINSKAYA STR., 61005, KHARKIV-5, UKRAINE
E-mail address: evtokarev@yahoo.com